



Jon Krumerman, M.D.

Cervical Fusion Surgery: Post-Operative Instructions

Incision Care

- Usually your stitches are beneath the surface of your skin. They will be absorbed into your body as you heal. Dermabond will be applied to incision site. It will come off naturally. Do not peel it off.
- You may shower, but no tub baths.
- You must keep your incision(s) dry for 7 days following your surgery.
- The bandage will need to be changed every day for 7 days. You will be given dressings for 7 days.
- The bandage may be removed and not replaced after 7 days. Do not apply anything to wound site. Keep clean and dry.

Pain

You may still have some pain, numbness or tingling in your neck or arm(s). This should decrease gradually as the nerves heal. An ice pack may be used on the incision to reduce swelling and pain. Heat may be used 24-48 hours after surgery for muscular discomfort.

Collar Instructions

- You should wear your neck collar while up only, unless other instructions are given at the time of your surgery. You should always wear neck collar while out or in a car.
- Hold your head straight while the collar is off. You may look up and down as well as rotate head 1 inch in all directions.
- Do not tilt head back to shampoo or shave.
- Your physician will determine when you are ready to wean out of your collar.
- You will have a bone growth stimulator to be worn 4 hours per day.

Call Our Office

If you...

- Feel severe pain, weakness or numbness in your leg(s)/arm(s).
- Notice excessive drainage, swelling or increased redness around your incision.
- Have a fever of 101 degrees or greater.
- Have a new problem controlling your bladder or bowel.
- Have difficulty breathing or calf pain.

Medication

- If you were given a Medrol Dosepak (steroid/anti-inflammatory medication) you may notice a slight increase in the above symptoms, as the Medrol Dosepak decreases. This is due to swelling and irritation around the nerve and will improve as you heal. Complete the full course of Medrol according to instructions in the package.
- You may be instructed to start taking a nonsteroidal anti-inflammatory medication (such as Ibuprofen, Naprosyn, Lodine) after you finish the Medrol Dosepak.
- You will be given a prescription for pain medicine when you are discharged from the hospital. Take as directed on the prescription. **Do not drive, operate machinery, or drink alcoholic beverages while taking this medication.**
- You may be given a prescription for muscle relaxants (to reduce spasms or tightness in your back). This medication may make you drowsy. **Do not drive, operate machinery, or drink alcoholic beverages while taking this medication.**

Medication (cont.)

- Medications will be refilled during normal business hours (8:30 am – 5:00 pm, M-Thrs, 9:30 am – Noon Fri), as the physician on call cannot renew prescriptions.
- Take a stool softener (i.e., Colace 100 mg twice daily) while on pain medications. If you do not have a bowel movement within 5 days of surgery and experience severe abdominal pain call your primary care physician or go to the E.R.

Activity

First Week:

- No auto travel. Once you get home from the hospital, stay there.
- Walk in your house only.
- Do not move your head more than 1 inch in all directions.
- Do not reach for objects above your head.
- Avoid house and yard work. You may cook, but do not lift more than 5 pounds until released to do so.
- No sexual activity.
- Do not drink from a bottle or can. Use a glass or a straw.
- Sleep on a firm mattress supporting your head on a pillow so it is in a level position.
- You may shower, but no tub baths if you have a hip incision. Pat dry, do not rub.
- Do not sleep with your arms over your head. You may lie on your back or side, but not on your abdomen.
- You may shampoo your hair in the shower only. Do not lean over the sink to shampoo. Hold your head in a neutral position.
- Do not cradle the phone between your cheek and shoulder.
- Take a nap or at least rest in the morning and again in the afternoon.
- No bending, lifting (greater than 10 lbs), twisting or straining.

Second Week:

- Start short walks outside, 1 to 2 blocks, twice a day. You may increase this as tolerated. Do not overdo it!
- Take a nap or at least rest in the morning and again in the afternoon.
- Continue to avoid house and yard work.
- You may ride as a passenger in a car, but do not drive.
- No sexual activity.
- No driving auto but may be a passenger.
- No bending, lifting (greater than 10 lbs), twisting or straining.

Third Week:

- You may start to drive short trips of 17 to 20 minutes. Make certain your seat, mirrors, and restraints are properly adjusted.
- You may resume sexual activity, do not support your body weight with your arms for the first few weeks.
- Increase walking distance as tolerated.
- You may be up and about as tolerated.
- Begin light housework. Do not sweep or vacuum. Stop for frequent breaks. Use good body mechanics.
- No bending, lifting (greater than 10 lbs), twisting or straining.

Activity (cont.)

Fourth Week:

- Gradually increase all activity as tolerated.
- Increase your walking distance to one mile a day, as tolerated.
- If approved by your physician, start isometric exercises as instructed.
- No bending, lifting (greater than 10 lbs), twisting or straining.

Follow-Up Appointment

Your first post-op appointment should be 2-3 weeks following your discharge from the hospital. Please call early to get a convenient time. You will be scheduled for an x-ray prior to your appointment.

Return to Work

Return to work varies with the type of occupation and will be discussed at your follow-up appointment.