



J. Michael Desaloms, M.D.

Cervical Fusion Surgery: Post-Operative Instructions

Incision Care

- Usually, your stitches are beneath the surface of the skin. They will be absorbed into your body as you heal. They are covered by Dermabond, a skin glue that protects your incision.
- You may shower but no tub baths. Avoid letting the shower spray on the incision. Pat incision dry after showering. Do not rub the Dermabond off.

Pain

You may still have some pain, numbness or tingling in your neck or arm(s). This should decrease gradually as the nerves heal. An ice pack may be used on the incision to reduce swelling and pain.

Collar Instructions

- You must wear your collar at all times.
- You may take it off to shower and shave.
- Hold your head straight while the collar is off.
- Do not tilt head back to shampoo or shave.
- Your physician will determine when you are ready to wean out of your collar.

Call Our Office

If you ...

- Feel severe pain, weakness or numbness in your leg(s)/ arm(s).
- Notice excessive drainage, swelling or increased redness around your incision.
- Have a fever of 101°F or greater.
- Have a new problem controlling your bladder or bowel.
- Have difficulty breathing or trouble swallowing.

Medication

- If you were given a Medrol Dosepak (steroid/ anti-inflammatory medication) you may notice a slight increase in the above symptoms, as the Medrol Dosepak decreases. This is due to swelling and irritation around the nerve and will improve as you heal. Complete the full course of Medrol according to the instructions in the package. Common side effects of steroids may include, but are not limited to, GI upset, insomnia and anxiety.
- You will be given a prescription for pain medicine when you are discharged from the hospital. Take as directed on the prescription. **Do not drive, operate machinery or drink alcoholic beverages while taking this medication.**
- You may have been given a prescription for muscle relaxants (to reduce spasms or tightness in your back). This medication may make you drowsy. **Do not drive, operate machinery or drink alcoholic beverages while taking this medication.**
- Medications will be refilled during normal business hours (8:30 am – 5:00 pm, Monday – Thursday and 8:30 am – 12:30 pm Friday) as the physician on call cannot renew prescriptions.



J. Michael Desaloms, M.D.

Cervical Fusion Surgery: Post-Operative Instructions (cont.)

Activity

First Week:

- No auto travel. Once you get home from the hospital, stay there.
- Walk in your house only.
- Do not move your head until released to do so.
- Do not reach for objects above your head.
- Avoid house and yard work. You may cook but do not lift more than 5 pounds until released to do so.
- Do not drink from a bottle or can. Use a glass or a straw.
- Sleep on a firm mattress supporting your head on a pillow so that it is in a level position.
- You may shower but no tub baths if you have a hip incision. Pat dry, do not rub.
- Do not sleep with your arms over your head. You may lie on your back or your side, but not on your abdomen.
- You may shampoo your hair in the shower only. Do not lean over the sink to shampoo. Hold your head in a neutral position.
- Do not cradle the phone between your cheek and shoulder.
- Take a nap or at least rest in the morning and again in the afternoon.

Second Week:

- Start short walks outside, 1 to 2 blocks, twice a day. You may increase this as tolerated. Do not overdo it!
- Take a nap or at least rest in the morning and again in the afternoon.
- Continue to avoid house and yard work.
- You may ride as a passenger in a car but do not drive.

Third Week:

- You may start to drive short trips of about 17 to 20 minutes. Make certain your seats, mirrors and restraints are appropriately adjusted.
- You may resume sexual activity, do not support your body weight with your arms for the first few weeks.
- Increase walking distance as tolerated.
- You may be up and about as tolerated.
- Begin light housework. Do not sweep or vacuum. Stop for frequent breaks. Use good body mechanics. After you have been cleared by the surgeon, you may lift up to 15 pounds.

Fourth Week:

- Gradually increase all activity as tolerated.
- Increase your walking distance to one mile a day, as tolerated.
- May walk on a treadmill, stationary bike or elliptical machine.



J. Michael Desaloms, M.D.

Cervical Fusion Surgery: Post-Operative Instructions (cont.)

Follow-Up Appointment

Your first post-op appointment should be 3-4 weeks following your discharge from hospital. Please call early to get a convenient time. You will be scheduled for an x-ray prior to your appointment.

Return to Work

Return to work varies with the type of occupation and will be discussed at your follow-up appointment.

Contact Numbers

Office: 214-750-3646

MA: Jessica Womble 214-445-6960

Appointments: Eve Freeman 214-445-6921