



Krishna Satyan, M.D.

Cervical Fusion Surgery: Post-Operative Instructions

Incision Care

- Usually, your stitches are beneath the surface of the skin. They will be absorbed into your body as you heal. They may be reinforced with steri-strips (sterile strips of tape) applied directly to the skin surface.
- Without Steri-strips, the dressing is a clear adhesive. It should start to peel off in a week or two.
- You may shower, but no tub baths, starting 3 days after surgery.
- You must keep your incision(s) dry for 2 days following your surgery.
- Steri-strips will come off on their own. Otherwise, will be removed at 1st post-op visit in 2 weeks.

Pain

You may still have some pain, numbness or tingling in your neck or arm(s). This should decrease gradually as the nerves heal. An ice pack may be used on the incision to reduce swelling and pain.

Collar Instructions

- You must wear your collar at all times when out of bed. Do not have to wear collar when sitting, showering or eating. Wear with activities, sleeping and car rides. No big bends when out of collar.
- You may take it off to shower and shave.
- Hold your head straight while the collar is off.
- Do not tilt head back to shampoo or shave.
- Your physician will determine when you are ready to wean out of your collar, but you are usually wearing the collar for 4 weeks.

Call Our Office

If you ...

- Feel severe pain, weakness or numbness in your leg(s)/ arm(s).
- Notice excessive drainage, swelling or increased redness around your incision.
- Have a fever of 101°F or greater.
- Have a new problem controlling your bladder or bowel.
- Have difficulty breathing.

Medication

- If you were given a Medrol Dosepak (steroid/ anti-inflammatory medication) you may notice a slight increase in the above symptoms, as the Medrol Dosepak decreases. This is due to swelling and irritation around the nerve and will improve as you heal. Complete the full course of Medrol according to the instructions in the package. Common side effects of steroids may include, but are not limited to, GI upset, insomnia and anxiety.
- You will be given a prescription for pain medicine when you are discharged from the hospital. Take as directed on the prescription. **Do not drive, operate machinery or drink alcoholic beverages while taking this medication.**



Krishna Satyan, M.D.

Cervical Fusion Surgery: Post-Operative Instructions (cont.)

- You may have been given a prescription for muscle relaxants (to reduce spasms or tightness in your back). This medication may make you drowsy. **Do not drive, operate machinery or drink alcoholic beverages while taking this medication.**
- Medications will be refilled during normal business hours (8:30 am – 5:00 pm, Monday – Thursday and 8:30 am – 12:30 pm Friday) as the physician on call cannot renew prescriptions.

The next morning, you may return home if you live out of town. You do not need to phone and check in but if you have questions, please do so.

IMPORTANT

The most dangerous but extremely rare risk of the same day disc surgery is the development of internal bleeding. This can put pressure on the spinal cord or nerves and cause paralysis. The potential crippling complication can be treated by reopening the incision and removing the blood, usually a clot. This is why the above precautions must be taken.

Activity

First Week:

- Limit auto travel. Passenger only. No driving
- No lifting > 10-20 lbs or big bending of neck for 3 months.
- Ambulate as much as tolerated. Avoid staying in one position as muscles will tighten.
- Sleep on a firm mattress supporting you head on a pillow so it is in a level position.
- You may shower but no tub baths. Pat the incision dry, do not rub.
- Do not sleep with your arms over your head. You may lie on your back or side, but no on you abdomen.
- You may shampoo your hair in the shower only. Do not lean over the sink to shampoo. Hold your head in a neutral position.
- Do not cradle the phone between your cheek and shoulder.
- May be fatigued for first few weeks. Rest/nap as needed.

Second Week:

- Increase walking distance as tolerated, working up to 1 mile per day
- May be fatigued for first few weeks. Rest/nap as needed.
- No driving until cleared at post-op visit.

Third Week:

- You may resume sexual activity, do not support your body weight with your arms for the first few weeks.
- Increase walking distance as tolerated, working up to 1 mile per day.
- You may be up and about as tolerated.
- Begin light housework. Do not sweep or vacuum. Stop for frequent breaks. Use good body mechanics.



Krishna Satyan, M.D.

Cervical Fusion Surgery: Post-Operative Instructions (cont.)

Fourth Week:

- Gradually increase all activity as tolerated.

Follow- Up Appointment

Your first post-op appointment should be 2-3 weeks following your discharge from the hospital. Please call early to get a convenient time. You will be scheduled for an x-ray prior to your appointment.